




Enhancing the potential effects of text messages delivered via an m-health intervention to improve packing of healthy school lunchboxes

Alison Brown^{1,2,3,4,*} , Rachel Sutherland^{1,2,3,4}, Lisa Janssen², Nayerra Hudson^{1,2}, Amelia Chooi^{1,2}, Renee Reynolds^{1,2}, Alison Walton⁵, Christophe Lecathelinais^{1,2,3,4}, Kathryn Reilly^{1,2,3,4}, Nicole Nathan^{1,2,3,4} and Luke Wolfenden^{1,2,3,4}

¹School of Medicine and Public Health, University of Newcastle, Callaghan, NSW, Australia: ²Hunter New England Population Health, Locked Bag 10, Wallsend, NSW 2287, Australia: ³Hunter Medical Research Institute, New Lambton Heights, NSW, Australia: ⁴Priority Research Centre for Health Behaviour, University of Newcastle, Callaghan, NSW, Australia: ⁵Mid North Coast Local Health District Health Promotion, Kempsey, NSW, Australia

Submitted 1 May 2020: Final revision received 20 September 2020: Accepted 2 October 2020: First published online 14 October 2020

Abstract

Objective: The aim of the study was to assess the impact of different lunchbox messages on parents' intention to pack a healthy lunchbox.

Design: This study employed an experimental design.

Setting: A series of messages were developed to align with the six constructs of the Health Belief Model. Messages were also developed that were (and were not) personalised and varied based on the source of the information provided (university, school, dietitian and health promotion service). During a telephone survey, participants were read the content of each message and asked about their intention to pack a healthy lunchbox.

Participants: Parents of primary school-aged children were randomised to receive different messages to encourage the packing of healthy lunchboxes.

Results: The study was completed by 511 parents. Linear mixed regression analyses identified significant differences ($P < 0.05$) in intention scores between variant messages targeting the same behavioural constructs for 'susceptibility', 'severity', 'benefits' and 'barriers' but not 'cues to action' or 'self-efficacy'. The highest mean behavioural intention score was for 'benefits', whilst the lowest mean score was for 'barriers'. There were no significant differences in intention scores of parents receiving messages from a dietitian, university, health promotion team or school ($P = 0.37$). Intention scores did not differ in which messages were personalised based on child's name ($P = 0.84$) or grade level ($P = 0.54$).

Conclusions: The findings suggest that messages that focus on the benefits of packing healthy lunchboxes may be particularly useful in improving intentions of parents to pack healthy foods for their children to consume at school.

Keywords

Text messaging
School lunchboxes
Parents
Children
Health Belief Model
Child diet
Message development

In countries such as Australia, the USA and the UK, the majority of school-aged children exceed dietary guideline recommendations regarding the consumption of discretionary foods, that is, those foods higher in salt, sugar and saturated fat^(1–3). Unhealthy dietary habits in children increase the risk of a variety of immediate and long-term health conditions, including dental caries, diabetes, obesity and CVD^(4–7). As such, improving child nutrition has been identified as a global health priority⁽⁸⁾.

Children consume approximately one-third of their daily energy intake during school hours⁽⁹⁾. Food consumed at school typically comes from either the school's food service, such as the school canteen or cafeteria, or via a packed lunch⁽⁹⁾. In Australia, like the UK⁽¹⁰⁾, the majority (>85%) of food consumed at school is brought from home via a school lunchbox, typically packed by parents⁽⁹⁾. In Australia, school lunchboxes have been found to contain an average of 3.5 serves of discretionary choices⁽¹¹⁾, far

*Corresponding author. Email alison.brown7@health.nsw.gov.au



exceeding Australian dietary recommendations for primary school-aged children⁽¹²⁾, and over 2800 kJ⁽¹³⁾, equivalent to 40 % of a primary school-aged child's entire daily energy intake⁽¹⁴⁾. As such, strategies that are successful in achieving even modest improvements in the nutritional quality of foods packed in school lunchboxes could make an important contribution to improving child nutrition⁽¹⁵⁾.

The use of text-based messaging via mobile phone applications has the potential to reach large numbers of parents and students at low cost and represents a promising intervention to improve the packing of healthy school lunchboxes⁽¹⁶⁾. Text message-based interventions have been found to be highly acceptable by end-users⁽¹⁶⁾, and systematic reviews have demonstrated that text-based messaging can be an effective tool for disease prevention and management across a variety of health behaviours and conditions⁽¹⁷⁾. Furthermore, a recent randomised trial utilising text messaging as part of a broader multi-component intervention found that it was effective in reducing discretionary choices and increasing 'everyday' healthy choices in school lunchboxes⁽¹⁸⁾.

Whilst there is evidence to suggest that text-based messaging may be effective in improving the nutritional quality of foods packed in a lunchbox, systematic reviews suggest that personalisation, tailoring of message content and message framing may enhance their effects⁽¹⁹⁾. Furthermore, the use of behaviour change frameworks in message development can help to identify factors that may improve intervention effects, and their use is recommended in the development of content for text message-based interventions⁽¹⁹⁾. The Health Belief Model (HBM) is one behaviour change theory that suggests that health risk behaviour is mediated by individual beliefs and perceptions^(20,21). The HBM has been used as a theoretical model in the development of mobile health or m-health interventions⁽²²⁾, and a number of its constructs have been found in meta-analyses to be effective in predicting a range of health behaviours⁽²³⁾. The HBM was designed to explain, and so help determine, which beliefs should be targeted in communication campaigns to modify health behaviours. As such, it may be a particularly relevant theory to apply in efforts to further enhance the impact of the text-message based strategies to improve lunchbox packing behaviours given the importance of communication for text-message based interventions.

Text-message based interventions have considerable potential reach⁽²⁴⁾ and may represent a lower cost means of delivering information at a population relative to in-person approaches. If delivered at a population level, even small improvements in their effect, achieved through deliberate and data-driven (optimisation)⁽²⁵⁾ approaches, may yield meaningful impacts at the population level. Therefore, the aim of this study was to assess:

1. The impact of different m-health message content targeting the same HBM construct on parents' intention to pack a healthy lunchbox.
2. The impact of m-health messages targeting different HBM constructs on parents' intention to pack a healthy lunchbox.
3. Whether the personalisation and credibility of the m-health message source increases parents' intention to pack a healthy lunchbox.

Method

Study design and setting

This study employed an experimental design. All participants were read one randomly selected message for each of the six domains of the HBM (six messages in total) and additionally one message which was (or was not) personalised and one message which varied based on the suggested source of the information provided (independent variable). After each message, participants were then asked about their intention to pack a healthy lunchbox (dependent variable). The study was conducted via a computerised-assisted telephone interview (CATI) of parents of primary school-aged children, from twelve Catholic primary schools in the Hunter New England region of New South Wales, Australia. The Hunter region encompasses major city and regional areas and is characterised by a high proportion of the population from low socioeconomic backgrounds⁽²⁶⁾.

Sample

The sampling frame for this study was parents with a child enrolled in Kindergarten to Grade 6 who had previously participated in an m-health randomised controlled trial of a physical activity and healthy lunchbox intervention⁽¹⁸⁾ and had consented to be contacted and invited to participate in future child health research studies.

Data collection and measures

Participants, who had consented to being contacted again in the m-health randomised controlled trial⁽¹⁸⁾, were telephoned to complete a survey via CATI between October 2018 and December 2018. Telephone interviews were delivered by experienced telephone interviewers.

Participant characteristics

Socio-demographic characteristics of parents and children assessed as part of telephone interviews conducted during the preceding m-health randomised controlled trial were used in this study, including parent postcode of residence, highest level of education, employment status and sex.

Message development

M-health messages were developed by a team of public health nutritionists, behavioural scientists and school education researchers. Content of messages was developed



to align with HBM constructs (susceptibility, severity, benefits, barriers, cues to action, self-efficacy)^(20,21). Within each HBM construct, a variety of messages were developed addressing topics relating to nutrition of children and lunchboxes, as described in Table 2:

1. Perceived susceptibility: defined as the belief that a person will develop a disease or condition⁽²⁷⁾. Five messages were developed targeting each of five common adverse effects of poor dietary intake reported in the literature to represent a concern for parents including child overweight and obesity⁽⁵⁾, Type II diabetes^(6,7), tooth decay⁽⁴⁾, dietary impacts on school performance and behaviour at school⁽²⁸⁾. All text messages used published estimates in Australian samples to communicate susceptibility of children to each of the adverse effects of poor child diet, such as the percent of children with overweight or obesity, tooth decay prevalence and risk of Type II diabetes.
2. Perceived severity: defined as the belief of the seriousness of a condition or the consequences associated with leaving a condition untreated⁽²⁷⁾. Four messages were developed targeting four of the most severe adverse effects of poor dietary intake reported in the literature including the effects of overweight and obesity^(5,28), Type II diabetes^(6,7) and tooth decay⁽⁴⁾. All text messages developed reported published statistics for the Australian context^(4,13,29) in an effort to communicate the seriousness and severity of conditions associated with poor dietary intake.
3. Perceived benefits: defined as the belief of potential positive aspects by partaking in a health action⁽²⁷⁾. Five messages were developed that highlighted five benefits of improved dietary intake for children, commonly reported in the literature and by formative evaluation with Australian parents including benefits of healthy eating at school and improved performance at school⁽²⁸⁾, formation of healthy habits throughout all life stages⁽¹²⁾, child tooth development⁽⁴⁾, impacts on mood and behaviour⁽²⁸⁾ and maintenance of a healthy weight⁽¹²⁾. Each topic was chosen to complement topics covered in messages aligned to the perceived susceptibility and severity constructs.
4. Perceived barriers: defined as the belief of the costs (both tangible and psychological) of partaking in a health action⁽²⁷⁾. Eight messages were created targeting the most common parental barriers in packing a healthy lunchbox based on formative evaluation with Australian parents in the current literature. Each message reflected one parental barrier including time constraints^(30–32), cost^(30–32), fussy eating^(30–32), food safety^(30,31), knowledge^(30–32), peer and parental influence⁽³¹⁾.
5. Cues to action: defined as factors or strategies that trigger action or readiness to change⁽²⁷⁾. Of the five messages developed, each message described simple strategies parents could use to improve the nutrition of school

- lunchboxes aligned to perceived parental barriers. The cues to action complemented the parental barriers of cost, food safety, knowledge and convenience^(30–32).
6. Self-efficacy: defined as the belief or confidence in a person's ability to take action⁽²⁷⁾. Five messages were developed that included 'how-to' information relating to parental barriers of packing a healthy lunchbox, in particular, cost, food safety, knowledge and skills, to complement cues to action messages^(30–32).

Additional messages were also developed to test the effects of different sources of information and personalisation of messages. Messages were developed varying the source of information as being from either an academic institution (the University of Newcastle) a local health service (Hunter New England Health, Health Promotion team), dietitians, or their local school, based on literature suggesting that educational systems, health services and health professionals are a trusted and credible source of information⁽³³⁾ as seen in Table 4. In addition, personalised messages were developed using the child's name and school grade level *v.* messages that did not use the child's name or grade level. Personalisation of messages using the participants or child's name or other socio-demographic features has been reported in the literature as being considered more meaningful and may lead to higher engagement with content⁽³⁴⁾. Examples of messages are highlighted in Table 4.

Instrument development and experimental manipulation

Each parent was randomly assigned one lunchbox message per HBM construct and one message each in relation to the credibility of the source and child's age and name personalisation. After interviewers read each message, participants were asked about their intention to pack a healthy lunchbox on a ten-point scale (0; no intention to pack a healthy lunchbox to 10; every intention to pack a healthy lunchbox). Intention was used as an indicator for packing a healthy lunchbox as it demonstrates a parent's reflective motivation to perform a behaviour or action, in this case, the packing a healthy lunchbox for their child⁽³⁵⁾. Survey questions were developed by a team of public health nutritionists, behavioural scientists and school education researchers based on similar items commonly used to assess behavioural intention^(35,36). Questions were pilot tested for comprehension and understanding and reviewed by experienced telephone interviewers.

Statistical analysis

All statistical analyses were performed using SAS (version 9.3) statistical software. Descriptive statistics were used to characterise the sample. Residential postcodes of parents were used to classify participants as residing in lower or higher socio-economic areas using the median score for

the state of New South Wales according to the 2016 Socio-Economic Index for Areas⁽³⁷⁾. Parents postcodes were also used to characterise the geographic locality of participants, as either rural or urban using the 2016 Accessibility/Remoteness Index of Australia⁽³⁷⁾. For each HBM construct, a linear mixed regression model was used to determine if there was a significant variation in behavioural intention scores between messages targeting that construct. Where significant effects ($P < 0.05$) were reported, using those same models, post hoc pair wise comparisons were undertaken to identify between which specific messages behavioural intention scores significantly differed. To assess differences in behavioural intention scores between each construct, a mean behavioural intention score for all messages in each construct was calculated, and linear mixed regression model was also used to compare these across constructs. Where significant effects ($P < 0.05$) were reported, post hoc pair wise comparisons were undertaken to identify between which constructs mean behavioural intention scores significantly differed. All models included a random effect for school to account for potential clustering. Statistical tests were two-tailed with an alpha of 0.05.

Results

Sample

From the original study, 790 (98%) parents agreed to be contacted for future studies. There were no significant differences on any socio-economic characteristics between parents who did and did not agree to be contacted. In total, 511 parents completed the CATI. Of the consenting and participating parents, 91% were female, 82% were from major cities, 85% were employed and 88% had an educational level of a certificate or diploma or higher. The characteristics of the consenting parents are listed in Table 1.

Table 1 Sample characteristics

Characteristics	n (%)	
Sex (n 499)		
Male	46	9
Female	453	91
Employment status (n 496)		
Employed	420	85
Unemployed	2	0.4
Domestic/home duties	57	11
Student	9	2
Other	8	2
Educational level (n 496)		
Some high school	27	5
Completed high school	41	8
Certificate or Diploma	163	33
University or college degree	265	53
Reside in areas of (n 510)		
Lower socio-economic status	269	53
Higher socio-economic status	241	47
Remoteness (n 510)		
Inner regional	92	18
Major cities	418	82

Impact of messages targeting the same Health Belief Model constructs on parents intention to pack a healthy lunchbox

Linear mixed regression analyses identified significant differences in behavioural intention scores between variant messages targeting the behavioural constructs of 'susceptibility', 'severity', 'benefits' and 'barriers' but not 'cues to action' or 'self-efficacy' (Table 2). Post hoc pair wise comparisons of the effects of individual messages within each construct revealed the following significant differences. For the HBM construct of 'susceptibility', messages reporting susceptibility statistics regarding tooth decay, overweight and obesity and poor classroom concentration linked to unhealthy food intake had higher behavioural intention scores than messages reporting statistics regarding discretionary foods ($P < 0.05$) typically packed in children's lunchboxes. Behavioural intention scores among parents receiving messages reporting susceptibility statistics on tooth decay were also significantly higher than those receiving messages regarding healthy foods typically packed in student lunchboxes ($P = 0.04$). For the HBM construct of 'severity', messages linking nutrition to diabetes risk had significantly higher behavioural intention scores than messages targeting overweight ($P = 0.02$). In relation to 'benefits' messages about protection against chronic disease scored higher messages regarding the benefits of good nutrition on academic performance ($P = 0.008$). For the HBM construct of 'barriers', the message reporting healthy lunchboxes often cost less than ones filled with unhealthy snacks had behavioural intention scores higher than messages suggesting that packing a lunchbox full of healthy everyday foods does not have to be time consuming ($P = 0.03$). Results relating to parent's intention to pack a healthy lunchbox based on the same HBM construct are highlighted in Table 2.

Impact of messages targeting different Health Belief Model constructs on parents intention to pack a healthy lunchbox

Linear mixed regression analyses identified significant differences in mean behavioural intention scores across HBM constructs ($P < 0.001$). The findings of post hoc pairwise comparisons are presented in Table 3. The highest mean behavioural intention score was for 'benefits' (8.57 (8.34; 8.80)), whilst the lowest mean score was 'barriers' (7.55 (7.32; 7.79)). 'Severity' and 'susceptibility' both scored significantly higher than 'barriers' (0.48, $P < 0.001$ and 0.63, $P < 0.001$), 'cues to action' (0.32, $P = 0.007$ and 0.47, $P < 0.001$) and 'self-efficacy' (0.41, $P < 0.001$ and 0.56, $P < 0.001$). There were no significant differences between 'barriers', 'cues to action' and 'self-efficacy', or 'cues to action' and 'self-efficacy' on behavioural intention scores.

**Table 2** M-health messages targeting the same Health Belief Model (HBM) constructs on parents' intention to pack a healthy lunchbox

	<i>n</i>	Mean score	95 % CI
HBM construct: susceptibility			
<i>Message 1</i> : 1 in 4 children in NSW are overweight, and healthy eating can help prevent this. ^A	95	8.51	8.02, 9.00
<i>Message 2</i> : 50 % of children in Australia have signs of tooth decay, and healthy eating can help prevent this. ^B	95	8.58	8.10, 9.07
<i>Message 3</i> : 40 % of children find it difficult to concentrate in class, and healthy eating can help prevent this. ^C	104	8.36	7.89, 8.82
<i>Message 4</i> : Only 12 % of primary school lunchboxes in the Hunter region contain only healthy foods. ^B	101	7.87	7.39, 8.34
<i>Message 5</i> : 86 % of primary school lunchboxes in the Hunter region contain unhealthy foods. ^{A-C}	108	7.65	7.19, 8.11
<i>P</i> value		0.02	
HBM construct: severity			
<i>Message 1</i> : Most children who eat sometimes foods at school find it harder to concentrate in class.	144	8.02	7.59, 8.46
<i>Message 2</i> : Poor nutrition can double your child's risk of developing Type II Diabetes. ^D	117	8.53	8.05, 9.01
<i>Message 3</i> : Almost half of 12-year-olds suffer from tooth decay, and nutrition plays an important role in this.	107	7.86	7.35, 8.37
<i>Message 4</i> : A quarter of primary school-aged children are above the healthy weight range, and the food your child consumes plays an important role in maintaining healthy growth. ^D	138	7.75	7.30, 8.19
<i>P</i> value		0.11	
HBM construct: benefits			
<i>Message 1</i> : Did you know that eating healthy food at school can help your child concentrate in the classroom and improve their academic performance? ^E	96	8.25	7.81, 8.69
<i>Message 2</i> : Did you know that eating well at school can help your child form healthy habits for life? ^F	96	8.33	7.89, 8.77
<i>Message 3</i> : Did you know that healthy eating plays an important role in looking after your child's teeth? ^F	110	8.54	8.13, 8.95
<i>Message 4</i> : Did you know that healthy eating is one of the best things you can do for your child's health and well-being? It improves their mood, behaviour and protects them against diseases like cancer and heart disease. ^E	108	9.08	8.67, 9.50
<i>Message 5</i> : Did you know that eating healthy food at school can help your child to maintain a healthy weight now and into the future?	93	8.51	8.06, 8.95
<i>P</i> value		0.06	
HBM construct: barriers			
<i>Message 1</i> : Are you short on time? Packing a lunchbox full of healthy everyday foods does not have to be time consuming. ^G	60	7.07	6.33, 7.80
<i>Message 2</i> : Lunchboxes filled with everyday healthy foods do not have to be expensive. In fact, healthy lunchboxes often cost less than ones filled with unhealthy snacks. ^G	47	8.32	7.49, 9.15
<i>Message 3</i> : Does your child dislike foods because of their colour, texture or the way they are prepared? Fussy eating is common, but parents are the perfect mentors to help children pack a lunchbox filled with everyday healthy foods.	65	7.51	6.80, 8.21
<i>Message 4</i> : Want to pack cold foods like yoghurt and cheese but worried about food safety? Follow these simple precautions to keep your lunchbox foods safe.	60	7.27	6.53, 8.00
<i>Message 5</i> : Packing an everyday lunchbox can be as easy as following these simple steps. Step 1: Pack veggies for crunch and sip; Step 2: Pack fruit and an everyday snack for recess; Step 3: Pack a healthy sandwich, roll, wrap or main item for lunch; Step 4: Finish with water as the perfect thirst quencher.	83	8.01	7.39, 8.64
<i>Message 6</i> : Do you have trouble finding a variety of healthy foods to pack in the lunchbox? Check out the SWAP IT sweet and savoury ideas for lots of great healthy inspiration that kids will love.	57	7.61	6.86, 8.37
<i>Message 7</i> : Do you have trouble with your children pestering you to pack unhealthy snacks in their lunchbox to be like other kids? Check out the SWAP IT page for ideas on how to tell your kids why it is important.	63	7.73	7.01, 8.45
<i>Message 8</i> : Wish more parents would pack more healthy lunchboxes? Do you feel pressure as a parent to pack the same foods as everyone else, even if they are unhealthy? Check out the SWAP IT page for some healthy inspiration on healthy foods kids love to pack in their lunchbox.	67	6.96	6.26, 7.65
<i>P</i> value		0.15	
HBM construct: cues to action			
<i>Message 1</i> : Need ideas on how to save money? Swap from a chocolate muffin to two pikelets and save money. Buy fruit and vegetables that are in season. Buy a big packet of rice crackers and make your own single serve portions.	117	7.87	7.36, 8.38
<i>Message 2</i> : Do not despair, SWAP IT is here to help with ideas on how to pack an everyday lunchbox for your fussy eater. Try the ¾ lunchbox rule: Pack 3 everyday items your child likes and 1 new everyday item each day.	83	7.64	7.03, 8.25
<i>Message 3</i> : An insulated lunchbox with an ice brick can keep the lunchbox 12 °C cooler. To keep the lunchbox extra cool on hot days, you could also freeze a water bottle or plain milk popper or yoghurt.	95	7.64	7.07, 8.21

Table 2 *Continued*

	<i>n</i>	Mean score	95 % CI
<i>Message 4:</i> Pack more fruit and veg into your child's day. Try vegetable sticks or fruit for Crunch&Sip®; include vegetable sticks or fruit and one other everyday snack, e.g. popcorn, yoghurt, rice crackers for recess; & pack a sandwich, wrap, roll or an alternative such as pasta salad full of veggies. On a scale of 0–10, how does this affect your intention to make a healthy swap?	90	7.78	7.19, 8.36
<i>Message 5:</i> There are loads of great pantry options that can go in the everyday Healthy lunchbox. Try: tinned fruit (in juice), tinned maize, baked beans, tuna; rice cakes or frozen berries.	114	7.57	7.05, 8.09
<i>P</i> value		0.93	
HBM construct: self-efficacy			
<i>Message 1:</i> Packing a lunchbox that does not blow the budget can be easy. Check out 'The Cost of Healthy Living' video to see the price difference between everyday and sometimes foods. Click the 'Visit Link' button for more money saving tips.	87	7.53	6.95, 8.11
<i>Message 2:</i> Packing a healthy lunchbox, even for the fussiest of eaters is simple. The SWAP IT programme has asked hundreds of kids about the everyday lunchbox foods they love to swap into their lunchbox. For more inspiration on packing an everyday lunchbox, click 'Visit Link' button or watch this video to find out what everyday foods kids love and why.	112	7.64	7.13, 8.16
<i>Message 3:</i> Keeping a lunchbox cold and free from foods that have spoiled is simple if you follow some easy steps. Click the 'Visit Link' button below for more great tips on keeping the lunchbox cold.	98	7.38	6.83, 7.93
<i>Message 4:</i> No need to be an expert chef or nutritionist to pack an everyday lunchbox that kids will love. For easy ideas on healthy lunchboxes, take a look at our Monday to Friday lunchbox packing video.	105	7.67	7.14, 8.20
<i>Message 5:</i> Every parent has the skills to pack a healthy lunchbox, download our 'Pantry Staples' PDF to take on your next shopping trip with lots of easy everyday healthy ideas.	96	7.84	7.29, 8.40
<i>P</i> value		0.81	

^{A-G}Similar subscripts delineate messages were significantly different to each other, at $P < 0.05$. Participants were asked about their intention to pack a healthy lunchbox on a ten-point scale (0; no intention to pack a healthy lunchbox to 10; every intention to pack a healthy lunchbox).

Table 3 Post hoc pairwise analyses comparing mean behavioural intention scores (to pack a healthy lunchbox) for Health Belief Model (HBM) constructs

HBM construct	<i>n</i>	Mean construct score	95 % CI	Post hoc between construct mean score difference	95 % CI
Susceptibility	504	8.19	7.96, 8.42	Susceptibility v.: Benefits: -0.38 Barriers: 0.63 Cues to action: 0.47 Self-efficacy: 0.56	-0.61, -0.15* 0.41, 0.86† 0.24, 0.70† 0.33, 0.79†
Severity	506	8.03	7.80, 8.27	Severity v.: Susceptibility: -0.15 Benefits: -0.53 Barriers: 0.48 Cues to action: 0.32 Self-efficacy: 0.41	-0.38, 0.07 -0.76, -0.31† 0.25, 0.71† 0.09, 0.54‡ 0.18, 0.63†
Benefits	507	8.57	8.34, 8.80	Benefits v.: Barriers: 1.01 Cues to action: 0.85 Self-efficacy: 0.94	0.79, 1.24† 0.62, 1.08† 0.71, 1.17†
Barriers	506	7.55	7.32, 7.79	Barriers v.: Cues to action: -0.17 Self-efficacy: -0.07	-0.39, 0.06 -0.30, 0.15
Cues to action	504	7.72	7.49, 7.95	Cues to action v.:	-0.14, 0.32
Self-efficacy	506	7.63	7.40, 7.86	Self-efficacy: 0.09	

*Significant difference to each other, at $P = 0.001$.

†Significant difference to each other, at $P < 0.001$.

‡Significant difference to each other, at $P = 0.007$.

Impact of personalisation and credibility of source of messages on parents' intention to pack a healthy lunchbox

There were no significant differences in behavioural intention scores of parents receiving messages from

a dietitian, a university, a health promotion team or the school ($P = 0.37$). Behavioural intention scores did not differ on items in which messages were personalised based on the child's name or grade level (Table 4).



Table 4 M-health messages targeting personalisation and credibility of source on parent's intention to pack a healthy lunchbox

	<i>n</i>	Mean score	95 % CI
How credible were the sources			
Dietitian	117	8.67	8.33, 9.00
University of Newcastle	121	8.88	8.55, 9.21
Hunter New England Health, Health Promotion Team	118	8.75	8.42, 9.09
School	141	8.50	8.19, 8.80
<i>P</i> value		0.37	
Personalised grade level			
Children in ^GRADERND^ find foods like fruit salad, cut-up veggies and crackers with hummus tasty. Try swapping these into your lunchbox and swapping out chips and muesli bars	250	7.66	7.29, 8.02
Children find foods like fruit salad, cut-up veggies and crackers with hummus tasty. Try swapping these into your lunchbox and swapping out chips and muesli bars.	246	7.80	7.44, 8.17
<i>P</i> value		0.54	
Personalised name			
Children like ^CHILDRND^ think foods like pikelets, scones and un-iced fruit buns are delicious. Try swapping these into your lunchbox and swapping out sweet biscuits, cakes and chocolates.	252	7.71	7.33, 8.09
Children think foods like pikelets, scones and un-iced fruit buns are delicious. Try swapping these into your lunchbox and swapping out sweet biscuits, cakes and chocolates.	244	7.66	7.28, 8.05
<i>P</i> value		0.84	

Discussion

The study found that the use of different HBM constructs in developing message content influenced parent's intention to pack a healthy lunchbox. Message content focussed on the benefits of packing healthy lunchboxes appeared to have particularly strong effects on behavioural intentions relative to messages related to other HBM constructs. Differences were also found in behavioural intention scores between messages assessing the same HBM construct. Personalisation of messages based on child name or school grade or altering the stated source of information provided in the message did not significantly influence parents' intention to pack a healthy lunchbox. The findings suggest that theoretically guided approaches in the development of text message-based intervention are likely to be effective in optimising the impact of such interventions in improving the foods consumed by children at school.

The study found that HBM constructs of 'benefits', 'susceptibility' and 'severity' were more influential than 'cues to action', 'self-efficacy' and 'barriers'. Lower behavioural intention scores on 'cues to action', 'self-efficacy' and 'barriers' suggest that such messages may have less salience.

The findings may reflect parent's belief that they are already packing a healthy lunchbox for their children or that they already have the skills and capacity to do so⁽³⁸⁾. If this is the case, incorporating a form of individualised feedback demonstrating the opportunity for improvement in packing of healthy lunchboxes, such as self-assessment tools, may improve the potential impact of messages targeting these constructs. Alternatively, the findings may reflect the limited ability of text messages aligned to these constructs (relative to others) to enhance behavioural intentions for the packing of healthy lunchboxes, suggesting that other strategies may be required.

Messages relating to benefits of healthy lunchboxes scoring higher than all other HBM constructs have been reported in other health promotion studies in parents. A study conducted in 2018 reported that messages relating to the reduction of sugar-sweetened beverages in children that were positive or 'gain-framed' had higher motivation scores in parenting practices than those that were negative or 'loss-framed'⁽³⁹⁾. Similarly, a meta-analysis conducted in 2014 which included 94 studies reported that gain-framed messages were significantly more likely to encourage health prevention behaviours than loss-framed messaging⁽⁴⁰⁾. The use of the HBM construct of benefits in framing messages may therefore highlight an impactful way to improve parents' intention to pack a healthy lunchbox.

Surprisingly, the personalisation of messages with children's names and grade levels had no effect on parents' intention to pack a healthy lunchbox, which has commonly been found in other studies^(17,19). A systematic review on the efficacy of text-based message interventions in health promotion conducted in 2013 reported that messages that were targeted and tailored were significantly associated with intervention efficacy with a greater effect size⁽¹⁹⁾. The review reported that tailoring of messages occurred in studies in a variety of ways including demographic and psychosocial tailoring⁽¹⁹⁾. The resulting no effect of personalisation of messages based on children's names or grade levels in our study may, therefore, be due to the level of tailoring and personalisation being too crude in only the name or grade level of the child changing in each message. More sophisticated approaches in the tailoring of lunchbox messages may provide better effects on behavioural intentions in the future. However, ethically obtaining and using personalised child information for the purpose of delivering health interventions may represent a considerable challenge for the application of such text-message based interventions. Similarly, the study found no effect on different information sources on intention to pack a healthy lunchbox. This may have been due to all sources of information included in the study considered to be credible⁽³³⁾.

The strengths of this study include the use of an experimental design in which parents were randomly allocated a series of messages. The use of an experimental design





strengthens the internal validity of the findings. Messages were also developed by an expert team of public health nutritionists, behavioural scientists and school education researchers using the current evidence base in relation to school lunchboxes and health behaviour risks. Nonetheless, the study had several limitations that must be considered when interpreting the results. Participants in the study were predominantly university educated from major cities. Additionally, the sampling frame was composed of parents who had previously participated in a physical activity and lunchbox intervention and may be more health oriented and motivated to improve health behaviour than the community at large. Such participant characteristics reduce the generalisability of the study findings. In addition, data on the ethnicity of parents were not collected which make it difficult to ascertain the representativeness of the sample. The use of a CATI in which parents were read out lunchbox messages instead of parents being able to visually see and read messages may have impacted on the validity of results. Differences exist in the attention, understanding and response to information presented in visual *v.* verbal formats^(41,42). As such, the effects of verbally delivered messages on behavioural intentions reported in this study may not generalise to delivery of the same message via text. Furthermore, in practice, the delivery of text-message based interventions also provides the opportunity to link users with other resources and information sources. This may be particularly important for HBM constructs such as barriers and cues to action where practical resources may be particularly helpful. Testing the effects of messages in a more naturalistic context, and in a form for which they may most optimally influence behaviour, may provide the most useful evidence to assess their effects. In future, recruiting parents from general community samples and testing the effects of messages with embedded links to further resources or information as appropriate may be a more applicable means of assessing their potential.

Conclusion

Notwithstanding the study limitations, the study provides useful insights that can assist in the development of text-based messages in an effort to improve the nutritional quality of school lunchboxes. Optimising the potential impact of messages in this way represents an efficient means of maximising the potential effects of such interventions prior to the conduct of large-scale randomised trials. Nonetheless, future randomised trials are required to confirm the effect of changes in message content on lunchbox packing behaviour of parents and dietary intake of children. Furthermore, a range of opportunities are available to enhance the impact of text message-based interventions beyond message content. For example, the timing and frequency of messages may influence its effects, as may

embedding other multimedia content or other strategies to enhance user experience and engagement^(19,43). The use of adaptive trial designs may be particularly useful in improving the efficiency and maximising the impact of text message-based interventions⁽⁴⁴⁾.

Acknowledgements

Acknowledgements: Not applicable. *Financial support:* The work was supported by Hunter Medical Research Institute (HMRI), Hunter Children's Research Foundation (HCRF) and Hunter New England Population Health. R.S. is supported by a National Health and Medical Research Council (NHMRC) TRIP Fellowship (grant number APP1150661); N.N. is supported by an NHMRC TRIP Fellowship (grant number APP1132450) and a Hunter New England Clinical Research Fellowship; LW is supported by an NHMRC Career Development Fellowship (grant number APP1128348), a Heart Foundation Future Leader Fellowship (grant number 101175) and a Hunter New England Clinical Research Fellowship. Funders had no role in the design, analysis or writing of this article. *Conflict of interest:* There are no conflicts of interest. *Authorship:* A.B., R.S., N.N. and L.W. conceived and designed the study. A.B., R.S., L.W., L.J., N.H., A.C., R.R., A.W. and K.R. designed the intervention. K.R. managed data collection. A.B. assisted with data cleaning. C.L. led statistical analysis. A.B. drafted the manuscript, with all other co-authors contributing to drafts of the paper. *Ethics of human subject participation:* This study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving research study participants were approved by the Hunter New England Human Research Ethics Committee (reference number 06/07/26/4.04), the University of Newcastle (reference number H-2008-0343) and the Maitland Newcastle Catholic Schools Office. Written informed consent was obtained from all subjects/patients.

References

1. Centre for Epidemiology and Evidence (2017) *What NSW Children Eat and Drink*. Sydney: NSW Ministry of Health.
2. Bates B, Cox L, Nicholson S *et al.* (2017) National Diet and Nutrition Survey. Results from Years 5–6 (combined) of the Rolling Programme (2012/13–2013/14) 2016; available at https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/551352/NDNS_Y5_6_UK_Main_Text.pdf (accessed December 2017).
3. U.S. Department of Health and Human Services and U.S. Department of Agriculture (2015) 2015–2020 Dietary Guidelines for Americans; available at <http://health.gov/dietaryguidelines/2015/guidelines/chapter-2/a-closer-look-at-current-intakes-and-recommended-shifts/> (accessed April 2020).



4. Armfield J, Spencer A & Brennan D (2009) *Dental Health of Australia's Teenagers and Pre-Teen Children: The Child Dental Health Survey, Australia 2003–04*. Canberra: AIHW, 2009. Contract No.: DEN 199. <https://www.aihw.gov.au/getmedia/0a8156ee-e7da-4ee2-be20-0244ba32c2bb/10778.pdf.aspx?inline=true> (accessed September 2019).
5. Australian Bureau of Statistics (2015) *Children's Risk Factors*. Canberra: Australian Bureau of Statistics.
6. Chiuvè SE, Sampson L & Willett WC (2011) The association between a nutritional quality index and risk of chronic disease. *Am J Prev Med* **40**, 505–513.
7. Must A, Spadano J, Coakley EH *et al.* (1999) The disease burden associated with overweight and obesity. *JAMA* **282**, 1523–1529.
8. World Health Organisation (2016) *Global Strategy on Diet, Physical Activity and Health: Childhood Overweight and Obesity*. Geneva: World Health Organisation; available at <http://www.who.int/dietphysicalactivity/childhood/en/> (accessed May 2016).
9. Bell AC & Swinburn BA (2004) What are the key food groups to target for preventing obesity and improving nutrition in schools? *Eur J Clin Nutr* **58**, 258–263.
10. Evans CE, Greenwood DC, Thomas JD *et al.* (2010) A cross-sectional survey of children's packed lunches in the UK: food- and nutrient-based results. *J Epidemiol Commun Health* **64**, 977–983.
11. Sanigorski AM, Bell AC, Kremer PJ *et al.* (2005) Lunchbox contents of Australian school children: room for improvement. *Eur J Clin Nutr* **59**, 1310–1316.
12. National Health and Medical Research Council (2013) *Australian Dietary Guidelines*. Canberra: National Health and Medical Research Council. Contract No.: 1864965754. https://www.eatforhealth.gov.au/sites/default/files/content/n55_australian_dietary_guidelines.pdf (accessed September 2019).
13. Sutherland R, Nathan N, Brown A *et al.* (2020) A cross-sectional study to determine the energy density and nutritional quality of primary-school children's lunchboxes. *Public Health Nutr* **23**, 1108–1116.
14. National Health and Medical Research Council (2006) *Nutrient Reference Values for Australia and New Zealand Including Recommended Dietary Intakes: Dietary Energy*. Australia: Commonwealth of Australia; available at <https://www.nrv.gov.au/dietary-energy> (accessed March 2020).
15. Cochrane T, Davey R & de Castella FR (2016) Estimates of the energy deficit required to reverse the trend in childhood obesity in Australian schoolchildren. *Aust N Z J Public Health* **40**, 62–67.
16. Payne HE & Lister C (2015) Behavioral functionality of mobile apps in health interventions: a systematic review of the literature. *JMIR Mhealth Uhealth* **3**, e20.
17. Cole-Lewis H & Kershaw T (2010) Text messaging as a tool for behavior change in disease prevention and management. *Epidemiol Rev* **32**, 56–69.
18. Sutherland R, Nathan N, Brown A *et al.* (2019) A randomized controlled trial to assess the potential efficacy, feasibility and acceptability of an m-health intervention targeting parents of school aged children to improve the nutritional quality of foods packed in the lunchbox 'SWAP IT'. *Int J Behav Nutr Phys Act* **16**, 54.
19. Head KJ, Noar SM, Iannarino NT *et al.* (2013) Efficacy of text messaging-based interventions for health promotion: a meta-analysis. *Soc Sci Med* **97**, 41–48.
20. Rosenstock IM (1974) Historical origins of the health belief model. *Health Educ Monogr* **2**, 328–335.
21. Becker MH (1974) The health belief model and personal health behavior. *Health Educ Monogr* **2**, 324–473.
22. Cho Y-M, Lee S, Islam SMS *et al.* (2018) Theories applied to m-health interventions for behavior change in low- and middle-income countries: a systematic review. *Telemed J E Health* **24**, 727–741.
23. Carpenter CJ (2010) A meta-analysis of the effectiveness of health belief model variables in predicting behavior. *Health Commun* **25**, 661–669.
24. Hall AK, Cole-Lewis H & Bernhardt JM (2015) Mobile text messaging for health: a systematic review of reviews. *Annu Rev Public Health* **36**, 393–415.
25. Wolfenden L, Bolsewicz K, Grady A *et al.* (2019) Optimisation: defining and exploring a concept to enhance the impact of public health initiatives. *Health Res Policy Syst* **17**. Published online: 30 December 2019. doi: 10.1186/s12961-019-0502-6.
26. Australian Bureau of Statistics (2012) *Population Estimates (HOIST)*. Sydney: NSW Ministry of Health.
27. Glanz K, Marcus Lewis F & Rimer BK (editors) (1997) *Theory at a Glance: A Guide for Health Promotion Practice*, 2nd ed. San Francisco: Jossey-Bass.
28. Carey FR, Singh GK, Brown HS 3rd *et al.* (2015) Educational outcomes associated with childhood obesity in the United States: cross-sectional results from the 2011–2012 National Survey of Children's Health. *Int J Behav Nutr Phys Act* **12**, Suppl. 1, S3.
29. Hardy LL, King L, Kelly B *et al.* (2010) Munch and move: evaluation of a preschool healthy eating and movement skill program. *Int J Behav Nutr Phys Act* **7**, 80.
30. Bathgate K & Begley A (2011) 'It's very hard to find what to put in the kid's lunch': what Perth parents think about food for school lunch boxes. *Nutr Diet* **68**, 21–26.
31. Hawthorne DL, Neilson IJ, Macaskill LA *et al.* (2018) Parental reports of lunch-packing behaviours lack accuracy: reported barriers and facilitators to packing school lunches. *Can J Diet Pract Res* **79**, 99–105.
32. Mazarello Paes V, Ong KK & Lakshman R (2015) Factors influencing obesogenic dietary intake in young children (0–6years): systematic review of qualitative evidence. *BMJ Open* **5**, e007396.
33. Dutta-Bergman M (2003) Trusted online sources of health information: differences in demographics, health beliefs, and health-information orientation. *J Med Internet Res* **5**, e21.
34. De Leon E, Fuentes LW & Cohen JE (2014) Characterizing periodic messaging interventions across health behaviors and media: systematic review. *J Med Internet Res* **16**, e93.
35. Michie S, van Stralen MM & West R (2011) The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* **6**, 42.
36. Francis J, Eccles MP, Johnston M *et al.* (2004) *Constructing Questionnaires Based on the Theory of Planned Behaviour: A Manual for Health Services Researchers*. Newcastle Upon Tyne, UK: Centre for Health Services Research, University of Newcastle Upon Tyne; available at <http://openaccess.city.ac.uk/id/eprint/1735> (accessed June 2019).
37. Australian Bureau of Statistics (ABS) (2001) *Technical Paper: Census of Population and Housing: Socio-Economic Indexes for Australia (SEIFA)*. Canberra: Commonwealth of Australia. <https://www.abs.gov.au/AUSSTATS/abs@nsf/allprimarymainfeatures/8C5F5BB699A0921CCA258259000BA619?open=document> (accessed June 2019).
38. Adamo KB & Brett KE (2014) Parental perceptions and childhood dietary quality. *Matern Child Health J* **18**, 978–995.
39. Zahid A & Reicks M (2018) Gain-framed messages were related to higher motivation scores for sugar-sweetened beverage parenting practices than loss-framed messages. *Nutrients* **10**, 625.
40. Gallagher KM & Updegraff JA (2012) Health message framing effects on attitudes, intentions, and behavior: a meta-analytic review. *Ann Behav Med* **43**, 101–116.



41. Kiat JE & Belli RF (2018) The role of individual differences in visual/verbal information processing preferences in visual\ verbal source monitoring. *J Cogn Psychol* **30**, 701–709.
42. Knoll AR, Otani H, Skeel RL *et al.* (2017) Learning style, judgements of learning, and learning of verbal and visual information. *Br J Psychol* **108**, 544–563.
43. Grady A, Yoong S, Sutherland R *et al.* (2018) Improving the public health impact of eHealth and mHealth interventions. *Austr N Z J Public Health* **42**, 118–119.
44. Pallmann P, Bedding AW, Choodari-Oskooei B *et al.* (2018) Adaptive designs in clinical trials: why use them, and how to run and report them. *BMC Med* **16**, 29.